



ISSN : 0973-7057

Int. Database Index: 616 www.mjl.clarivate.com

Antagonistic activity of lactic acid bacteria isolated from human colostrum against common human pathogens

Komalben J. Hirani^{a*}, Amar P. Garg^a & Sandeep K. Srivastava^b

^aSchool of Biological Engineering & Life Sciences, Shobhit Institute of Engineering & Technology, Deemed to-Be-University, Modipuram, Meerut, India

^bCentre for Innovation, Research and Development, Dr. B. Lal Institute of Biotechnology, Jaipur, India.

Received : 16th July, 2022 ; Revised : 16th August, 2022

Abstract- The present study was carried out to isolate those probiotic strains of lactic acid bacteria (LAB) from human colostrum that possess antibacterial activity against wide range of human pathogens. All the isolates were biochemically characterized and studied for their probiotic properties including acid tolerance, production of bacteriocin and hydrogen peroxide, tolerance to different concentrations of bile salts and resistance against common antibiotics. Antibacterial activity of the isolated species was tested against different Gram's negative as well as positive pathogens including *Klebsiella pneumoniae*, *Proteus vulgaris*, *Staphylococcus aureus*, *Salmonella typhi*, *Escherichia coli* and *Pseudomonas aeruginosa*. All isolates showed inhibitory effects against above mentioned bacteria. L15002 showed high antibacterial activity against all test pathogens. It was genotypically confirmed as *Lactococcus lactis*. The growth of bacteria was assessed *in vitro* using viable bacterial counting. Antibacterial response of *Lactococcus lactis* was attributed to the production of bacteriocins which were responsible for cell death as well as cell inhibition. The present study is focused on antagonistic activity of different species of LAB isolated from human colostrum.

Key words: Lactic acid bacteria, Probiotics, Bacteriocins, Antagonistic activity

INTRODUCTION

Human milk is the gold standard for infant nutrition as it contains multiple biologically active components, including immune factors, lipids, oligosaccharides, microRNAs hormones and various other nutrients in different quantities.^{1,2} It promotes the infant growth and development, and WHO has recommended exclusive breast feeding during the first 6 months of life.³ Microorganisms have emerged as important bioactive components of human milk that play an important role as commensal, mutualistic

or potentially probiotic bacteria in healthy human milk,⁴ which has created increasing interest in human milk microbiology and their effects in maternal-infant health. Species of *Lactobacillus salivarius*, *Lactobacillus fermentum*, *Lactobacillus gasseri*, *Bifidobacterium breve*, *Bifidobacterium adolescentis* and *Bifidobacterium longum* sub sp. *infantis* have demonstrated potential to promote mother and infant health, including the prevention or treatment of lactational mastitis.^{5,6}

Antibacterial effects of lactic acid bacteria (LAB) against human pathogens are well documented in scientific literature where these have been isolated from different

*Corresponding author :

Phone : 7874119409

E-mail : komalhirani2008@gmail.com

milk laying animals such as buffalo, cow, goat and camel milk.⁷ Previous studies have reported the antagonistic activity of *Enterococcus* spp., *Pediococcus* spp., *Lactobacillus* spp and species of *Lactococcus*.⁸ LAB have high potential of probiotic properties that are generally acceptable for human consumption.⁹ LAB have high industrial importance in fermentation and pharmaceutical industries due to their wide range of applications.¹⁰ They play a significant role in developing immune system of infants.¹¹ Large number of bacteria with high probiotic potentials enter infant's gut through breast feeding in initial days of life¹² that protects the infant immune system in fighting against wide variety of pathogens.¹³ They are very commonly reported in the gut of adults.¹⁴ Problems related indigestion are attributed to the imbalance of gut microbiota.¹⁵ LAB produce large amounts of other inhibitory substances like different bacteriocins, lactocins and some other organic acids,¹⁶⁻¹⁹ inhibit the growth of other microorganisms. Mechanism of microbial antagonism is attributed to the competition for nutrients, space, secretion of inhibitory substances, stimulation of immune system and adhesion inhibition of pathogens to mucosal surface and various other factors.^{13,20} The rapid increase in emergence of chronic toxicity caused through drug-resistant strains has encouraged the researchers for developing alternate treatment for controlling and treating bacterial infections.^{21,22} This paper deals with the antagonistic activity of LAB against common human pathogens with the objective to find out their probiotic potential.

MATERIALS & METHODS

Colostrum samples were collected from the maternity ward of JNU hospital, Jaipur after approval from institutional ethics committee. Samples were serially diluted in sterile peptone water and 0.1mL of inoculum was aseptically inoculated on MRS (deMan, Rogosa & Sharpe) agar plates. The plates were incubated anaerobically at 37°C for 24-48h. Isolates were sub-cultured to get pure colonies and were identified phenotypically using different biochemical tests. Identification and biochemical tests were carried out at Shobhit Institute of Engineering & Technology, Meerut.

Probiotic properties of isolates were checked using different tests as previously mentioned in (Ritesh *et al.*, 2020 a, b, c)²³⁻²⁵.

Test organisms

All six strains of pathogens used in the present study are already available in our laboratories. The indicator pathogens used in the study were *Staphylococcus aureus* ATCC25923, *Proteus vulgaris* ATCC33420, *Escherichia coli* ATCC25922, *Klebsiella pneumonia* MTCC3384, *Salmonella typhi* MTCC733 and *Pseudomonas aeruginosa* ATCC27853. All six pathogens were overnight grown in nutrient broth at 37°C.

Anti-bacterial activity of LAB


The antagonistic activity of isolates against test pathogens was measured using agar well diffusion method. The pathogens were inoculated on Mueller Hilton agar plates and with the help of sterile cork borer (6 mm in diameter), agar surface was punctured to create wells. Liquid cultures of isolates grown overnight in MRS broth were poured in the wells with the help of micropipette. The plates were allowed to incubate at 37°C for 24-48h in upright position. Zones of inhibition were observed and measured using zone measuring disc after completion of incubation.^{26,27}

RESULTS & DISCUSSION

Eight isolates which were showing best probiotic properties were screened for testing the antagonistic activity against common human pathogens. These isolates were morphologically and biochemically identified using Gram's staining, catalase test, oxidase and sugar fermentation tests as recommended in Bergey's Manual of Systematic Bacteriology and earlier described by Bisht and Garg (2019)²⁷. All the 8 isolates were Gram's positive, catalase and oxidase negative in nature. On the basis of their ability to ferment different sugars, genus of the isolates was identified that included 3 isolates of *Lactobacillus*, 3 of *Lactococcus* and 2 were *Pediococcus*. All the eight isolates were tolerant to acidic pH, and bile salts and were also resistant to different antibiotics. Antagonistic test revealed that isolate (I4601) was showed highest antibacterial activity against *Staphylococcus aureus* ATCC25923 (Fig.1), *Proteus vulgaris* ATCC33420, *Escherichia coli* ATCC25922 (Fig.1), *Klebsiella pneumonia* MTCC3384 while *Salmonella typhi* MTCC733 *Pseudomonas aeruginosa* ATCC27853 were inhibited by isolate no I0502. The isolate (I4601) isolate was identified as *Lactococcus lactis* (Table1).

Table 1: Zone of inhibition of selected lactic acid bacteria

Isolate Name	<i>Staphylococcus aureus</i> ATCC25923	<i>Proteus vulgaris</i> ATCC33420	<i>Escherichia coli</i> ATCC25922	<i>Klebsiella pneumoniae</i> MTCC3384	<i>Salmonella typhi</i> MTCC733	<i>Pseudomonas aeruginosa</i> ATCC27853
Zone of Inhibition (mm)						
I2501	16mm	5mm	8mm	5mm	8mm	10mm
I3302	19mm	6mm	10mm	04mm	09mm	10mm
I4601	24mm	8mm	12mm	09mm	16mm	13mm
I0502	21mm	7mm	14mm	06mm	18mm	18mm
I5201	21mm	8mm	10mm	04mm	14mm	16mm
I5601	23mm	6mm	06mm	07mm	10mm	10mm
I0601	20mm	7mm	09mm	06mm	06mm	09mm
I5502	18mm	5mm	07mm	05mm	06mm	08mm

*S. aureus* ATCC25923*E. coli* ATCC25922**Fig 1: Antagonistic activity of *Lactococcus lactis* against *Staphylococcus aureus* and *Escherichia coli***

The scientific interest in studies on human milk bacteria is mainly focused on (a) the roles that milk bacterial communities play on infant health and development, (b) their influence on the colonization of the gut microbiota in early life; (c) the roles that they may play on maternal health, including breast health; (d) the maternal, infant, environmental and medical factors that shape and modulate their compositions, (e) their antagonistic activity with different enteric pathogens (f) origin, in order to confirm or refuse the existence of an endogenous oral-entero-mammary route allowing the selective translocation of some bacteria from the maternal digestive tract to the mammary gland, (h) differential microflora of normal or scissor deliveries and (i) impact of diet on composition of microflora of human colostrum.

The present investigation aimed to isolate and characterize human colostrum bacteria and to find out their antagonistic activity against common human pathogens. We identified 3 isolates of *Lactobacillus*, 3 of *Lactococcus* and 2 of *Pediococcus* on MRS agar medium under anaerobic/partially aerobic conditions. In view of the

antagonistic activity of LAB and its commercial applications in food industry,²⁸⁻³¹ the microbiologists are isolating potentially antagonistic LABs from different sources and evaluating their antimicrobial activities against common food borne human pathogens. Almedia *et al.*, (2021)³² have recently characterized a large number of functional and taxonomic groups of human gut microbiota and have presented a Unified Human Gastrointestinal Genome (UHGG) collection comprising 204,938 non-redundant genomes from 4,644 prokaryotes which encode >170 million proteins and have highlighted the importance of studies on gut microbiome. Roughly 40 trillion microbes, consisting of more than 3000 species of several viruses, bacteria, archaea, fungi and other eukaryotes live in and on our body and are extremely important for our health, however, few cause diseases and disorders.³³ Microbial biomass in/on our body consists of 1 kg (approximately) representing about 1,000,000 genes while entire human genome consists of approximate 30,000 genes and the human microbial genome greatly influence the physiology and behaviour of human. Gut microbiome is the principal ecological niche and secretes various enzymes that are mainly responsible for the digestion of food and also accounts for 70% of the total immunity of the body. Lactic acid bacteria (LAB) constitute the major part of gut microbiome but these are difficult to culture because of the requirement of anaerobic / partially aerobic conditions for their culture. Most scientists have focused on genomic characterization of gut microbiota and a large number of microbial species have been identified,³⁴ however, very few species have been cultured so far, mainly because of the problems of quick death of gut microbes in between the sampling and processing. The

isolation, characterization and culture of 8 isolates of LAB from human colostrum as representative of gut microbiota of human in this paper suggest that gut microbes if cultured under anaerobic / partially aerobic conditions on MRS agar media, can help in isolation of more and more species which can then be evaluated for their commercial exploitation. Diet influences the gut microbiome³⁵ and the importance of plant-based diet in regulation of gut microbiome and its impact on human brain functions has been beautifully reviewed by Medawar *et al.* (2019)³⁶. Good bacteria can also help to manage depression.³⁷ It is suggested that human gut micro biome need to be cultured so that their probiotic value may be evaluated and used for commercial purposes. The presence of good bacteria in human colostrum having antagonistic activity suggest that breast feeding to infants provide protection to infants from enteric pathogens and the recommendations of WHO for initial 6-month breast feeding have scientific value and advantage in growth, development and protection of child against infections.

CONCLUSIONS

LAB have great potential of fighting against human pathogens. Consumption of LAB rich dairy products and fermented foods may greatly benefit our digestive and immune system including gut microbiome. Keeping in view the limitations of *in vitro* studies, it is suggested that further *in vivo* studies are required for clear picture of the role of LABs. Better strains of LAB as probiotics are required to solve the problems related to deficiency of probiotics.

REFERENCES

1. **Rodríguez J. M.** 2014. The origin of human milk bacteria: is there a bacterial entero-mammary pathway during late pregnancy and lactation? *Adv. Nutr.* **5**:779-784
2. **Hennet T. & Borsig L.** 2016. Breastfed at tiffany's. *Trends Biochem. Sci.* **41**: 508–518.
3. **World Health Organization.** 2003. Global Strategy for Infant and Young Child Feeding. Geneva: WHO.
4. **Fernández Leónides, Susana Langa, Virginia Martín, Antonio Maldonado, Esther Jiménez, Rocío Martín, and Juan M. Rodríguez.** 2013. The human milk microbiota: origin and potential roles in health and disease. *Pharmacol.Res.* **69**: 1-10.
5. **Fernández Leónides, Nivia Cárdenas, Rebeca Arroyo, Susana Manzano, Esther Jiménez, Virginia Martín, and Juan Miguel Rodríguez** 2016. Prevention of infectious mastitis by oral administration of *Lactobacillus salivarius* PS2 during late pregnancy. *Clin. Infect. Dis.* **62**: 568–573.
6. **Shin Suk Pyo, Yoon Mi Choi, Won Hee Kim, Sung Pyo Hong, Jong-Min Park, Joohee Kim, Oran Kwon, Eun Hyun Lee, and Ki Baik Hahm.** 2018. A double blind, placebo-controlled, randomized clinical trial that breast milk derived- *Lactobacillus gasseri* BNR17 mitigated diarrhea-dominant irritable bowel syndrome. *J. Clin. Biochem. Nutr.* **62**:179–186.
7. **Caggianiello G., Kleerebezem M. and Spano G.** 2016. Exopolysaccharides produced by lactic acid bacteria: from health-promoting benefits to stress tolerance mechanisms. *Appl. Microbiol. Biotechnol.* **100**:3877–3886.
8. **Liu Zhengqi, Zhihong Zhang, Liang Qiu, Fen Zhang, Xiongpeng Xu, Hua Wei, and Xueying Tao.** 2017. Characterization and bioactivities of the exopolysaccharide from a probiotic strain of *Lactobacillus plantarum* WLPL04. *Journal of Dairy Science.* **100(9)**: 6895-6905.
9. **Dertli E., Mayer M. J. and Narbad A.** 2015. Impact of the exopolysaccharide layer on biofilms, adhesion and resistance to stress in *Lactobacillus johnsonii* FI9785. *BMC Microbiol.* **15(1)**:1-9.
10. **Nataraj B. H., Ali S. A., Behare P. V. and Yadav H.** 2020. Postbiotics-parabiotics: the new horizons in microbial biotherapy and functional foods. *Microb. CellFact.* **19(1)**:1-22.
11. **Fanning S., Hall L. J., Cronin M., Zomer A., MacSharry J., Goulding D., O'Connell Motherway M., Shanahan F., Nally K., Dougan G. and van Sinderen D.** 2012. Bifidobacterial surface-exopolysaccharide facilitates commensal-host interaction through immune modulation and pathogen protection. *Proceedings of the National Academy of Sciences.* **109(6)**: 2108-2113.

12. Saadat R. Y., Khosroushahi Y. A., and Gargari P. B. 2019. A comprehensive review of anticancer, immune modulatory and health beneficial effects of the lactic acid bacteria exopolysaccharides. *Carbohydr. Polym.* **217**: 79–89.
13. Walker W.A. and Iyengar R. S. 2014. Breast milk, microbiota, and intestinal immune homeostasis. *Pediatr. Res.* **77**: 220–228.
14. Matsubara V. H., Wang Y., Bandara H., Mayer M. P. A. and Samaranayake L. P. 2016. Probiotic lactobacilli inhibit early stages of *Candida albicans* biofilm development by reducing their growth, cell adhesion, and filamentation. *Appl. Microbiol. Biotechnol.* **100**: 6415–6426.
15. Lebeer S., Vanderleyden J., and De Keersmaecker S. C. J. 2010. Host interactions of probiotic bacterial surface molecules: comparison with commensals and pathogens. *Nat. Rev. Microbiol.* **8**: 171–184.
16. Kanmani P., Suganya K., Yuvaraj N., Pattukumar V., Paari K. A., & Arul V. 2013. Synthesis and functional characterization of antibiofilm exopolysaccharide produced by *Enterococcus faecium* MC13 isolated from the gut of fish. *Applied biochemistry and biotechnology.* **169**(3): 1001-1015.
17. Jeong Dana, Dong-Hyeon Kim, Il-Byeong Kang, Hyunsook Kim, Kwang-Young Song, Hong-Seok Kim, and Kun-Ho Seo. 2017. Characterization and antibacterial activity of a novel exopolysaccharide produced by *Lactobacillus kefiranoferiens* DN1 isolated from kefir. *Food Control.* **78**: 436-442.
18. Graf K., Last A., Gratz R., Allert S., Linde S., Westermann M., Gröger M., Mosig A. S., Gresnigt M. S. and Hube B. 2019. Keeping *Candida* commensal: how lactobacilli antagonize pathogenicity of *Candida albicans* in an in vitro gut model. *Disease models & mechanisms*, **12**(9): p.dmm 039719.
19. Olaimat A. N., Aolymat I., Al-Holy M., Ayyash M., Abu Ghoush M., Al-Nabulsi A. A., Osaili T., Apostolopoulos V., Liu S. Q. and Shah N. P. 2020. The potential application of probiotics and prebiotics for the prevention and treatment of COVID-19. *npj Science of Food*, **4**(1):1-7.
20. Mahdhi Abdelkarim, Nadia Leban, Ibtissem Chakroun, Mohamed Aymen Chaouch, Jawhar Hafsa, Kais Fdhila, Kacem Mahdouani, and Hatem Majdoub. 2017. Extracellular polysaccharide derived from potential probiotic strain with antioxidant and antibacterial activities as a prebiotic agent to control pathogenic bacterial biofilm formation. *Microbial pathogenesis.* **109**: 214-220.
21. Xu Yuanmei, Yanlong Cui, Fangfang Yue, Lihua Liu, Yuanyuan Shan, Bianfang Liu, Yuan Zhou, and Xin Lü. 2019. Exopolysaccharides produced by lactic acid bacteria and Bifidobacteria: Structures, physiochemical functions and applications in the food industry. *Food Hydrocolloids.* **94**: 475-499.
22. Xu Xiaoqing, Qing Peng, Yuwei Zhang, Dandan Tian, Pengbo Zhang, Ying Huang, Lan Ma, Vermont P. Dia, Yu Qiao, and Bo Shi. 2020. Antibacterial potential of a novel *Lactobacillus casei* strain isolated from Chinese northeast sauerkraut and the antibiofilm activity of its exopolysaccharides. *Food & Function.* **11**(5): 4697-4706.
23. Ritesh Kumar Arya, Jaspreet Singh and Amar Garg. 2020 a. Isolation, Characterization and Role of Lactic Acid Bacteria in Human Colostrum. *Biosci. Biotech. Res. Comm.* **13**(1): 295–306.
24. Ritesh Kumar Arya, Jaspreet Singh and Amar Garg. 2020 b. Detection and Enumeration of Lactic Acid Bacteria Isolated from Human Colostrum using Traditional Microbiology Techniques. *Advances in Bioresearch.* **11**(4): 84–88.
25. Ritesh Kumar Arya, Jaspreet Singh and Amar Garg. 2020 c. In vitro Evaluation of Probiotic Potentials of Lactic Acid Bacteria Isolated from Human Colostrum. *Advances in Bioresearch.* **11**(4): 93–99.
26. Balouiri M., Sadiki M. and Ibnsouda S. K. 2016. Methods for in vitro evaluating antimicrobial activity: are view. *J. Pharmaceut. Anal.* **6**: 71-79.
27. Bisht N. and A. P. Garg. 2019. Isolation, characterization and probiotic value of lactic acid bacteria from milk and milk products. *Biotech. Today.* **9**(2): 54-63

28. Aoudia Nabil, Aurélie Rieu, Romain Briandet, Julien Deschamps, Johanna Chluba, Gaëtan Jegou, Carmen Garrido, and Jean Guzzo. 2016. Biofilms of *Lactobacillus plantarum* and *Lactobacillus fermentum*: Effect on stress responses, antagonistic effects on pathogen growth and immunomodulatory properties. *Food Microbiology*. **53**: 51-59.
29. Bhat B., and Bajaj B. K. 2018. Hypocholesterolemic and bioactive potential of exopolysaccharide from a probiotic *Enterococcus faecium* K1 isolated from kalarei. *Bioresour. Technol.* **254**:264–267.
30. Fernández L., Ruiz L., Jara J., Orgaz B. and Rodríguez J. M. 2018. Strategies for the preservation, restoration and modulation of the human milk microbiota. Implications for human milk banks and neonatal intensive care units. *Front Microbiol.* **9**:2676.
31. Ayyash Mutamed, Basim Abu-Jdayil, Pariyaporn Itsaranuwat, Emmanuel Galiwango, Camila Tamiello-Rosa, Hassan Abdullah, Gennaro Esposito, Yamanappa Hunashal, Reyad S. Obaid, and Fathalla Hamed. 2020. Characterization, bioactivities, and rheological properties of exopolysaccharide produced by novel probiotic *Lactobacillus plantarum* C70 isolated from camel milk. *International Journal of Biological Macromolecules*. **144**: 938-946.
32. Almedia Alexandre, Stephen Nayfach, Miguel Boland. 2021. *Nature Biotechnology*. **23**:105-114
33. Sundararaman A., Ray M., Ravindra P., and Halami P. M. 2020. Role of probiotics to combat viral infections with emphasis on COVID-19. *Appl. Microbiol. Biotechnol.* **104**: 8089–8104.
34. Samuel C. Forster, Nitin Kumar, Blessing O, Alexandre Elisa Viciani. 2019. A human gut bacterial genome and culture collection for improved metagenomic analyses. *Nature Biotechnology*. **37**: 186-192
35. Angelis MariaDe, Ilario Ferrocino, Francesco Maria Calabrese, Francesca DeFilippis, F., Cavallo, N. and Siragusa S. 2020. Diet influences the functions of the human intestinal microbiome. *Scientific Reports*. **10**(1):42-47
36. Medawar Evelyn, Sebastian Huhn, Arno Villringer and Veronica Witte 2019. The effects of plant-based diets on the body and the brain: a systematic review. *Translational Psychiatry*. **9**: 226.
37. Chevalier G., Siopi E., Guenin-Macé L., Pascal M., Laval T., Rifflet A., Boneca I. G., Demangel C., Colsch B., Pruvost A. and Chu-Van E. 2020. Effect of gut microbiota on depressive-like behaviors in mice is mediated by the endocannabinoid system. *Nature Communications*. **11**(1):1-15.
