

ISSN: 0973-7057

A nematode genus *Eustrongylides* Jagerskiold, 1909 from Karbhala wetland, Cachar District, Assam, first host report and geographical report

Khaidem Binkya, M. Shomorendrab* & Devashish Karc

^aDepartment of Zoology, Pravabati College, Mayang, Imphal, Manipur, India ^bDepartment of Zoology, Thambal Marik College, Oinam, Manipur, India ^cDivision of Wetlands, Fishery Science and Aquaculture, Department of Life Sciences and Bioinformatics, Assam University, Silchar, Assam, India

> Received: 25th November, 2024; Revised: 04th January, 2025 DOI:-https://doi.org/10.5281/zenodo.15978195

Abstract- In the present study, in order to extend the knowledge on the distribution of parasites of fish of the genus *Eustrongylides* in the area were undertaken. *Eustrongylides* is a genus of nematode belonging to the family Dioctophymidae. Fish specimens were collected from Karbhala wetland and analysed. Larval form of *Eustrongylides* sp. were found from the air bladder and intestine of fish host *Channa punctata*. The presence of the parasite in the local fresh water fish is currently very low. The present paper deals with the first report and geographical report of a nematode belonging to the genus *Eustrongylides* from the fish host *Channa punctata* from Karbhala wetland, Cachar District, Assam. Future investigations will show if the parasite expands in these areas. The taxonomical observation was briefly discussed.

Keywords: Nematode genus Eustrongylides, Kharbala wetland, new host record & new locality

INTRODUCTION

In 1909, Jagerskiold erected the genus Eustrongylides with E. tubifex as its type species from Anas, Mergus, Pelecanus, Colymbus, Podiceps, Alca etc.¹ The only species representative reported from fish was E. wenrichi Canavan, 1929.² In 1963, Fernando and Furtado were reported the larvae of this genus from three fishes, viz., Heteropneustes fossilis, Ompok bimaculatus and Wallago attu from Sri Lanka.³ In 1974, Kalyankar reported another larvae of the genus from Sperata seenghala from Nanded, Maharashtra.⁴ In 1979 Naidu reported from Mastacembelus armatus from Maharashtra.⁵ In 1997, Wang et al. reported the larvae of Eustrongylides africanus in Clarias from Bida flood plain in Nigeria from the fish

*Corresponding author: Phone: 9862490384

E-mail: smaibam90@gmail.com

species of the families Engraulidae, Cyprinidae, Siluridae, Bagridae, Channidae and Percichthyidae. In 2004, 2005, 2008, Ibiwoye *et al.* studied and reported the occurrence of *Eustrongylides africanus* larvae in *Clarias* species. Percies Perci

MATERIALS & METHODS

Fishes of different sizes were routinely collected from Karbhala wetland, Cachar, Assam. The fishes were examined for endo and exo parasitic infection. The worms collected from the fishes were preserved in 70% alcohol. The preserved specimens were cleared in lactophenol and observed under microscope. Photographs were taken using a digital camera. Drawing was done using mirror type Camera Lucida. The measurements were given in millimetres. Identification was done through standard literature.

Biospectra: Vol. 20(1), March, 2025

An International Biannual Refereed Journal of Life Sciences

RESULT & DISCUSSION

Eustrongylides sp. (Figure 1, Plate 1)

Family - Dioctophymatidae, Railliet, 1915 Genus - *Eustrongylides* Jagerskiold, 1909

Description- Body 14.1 - 16.92 mm in length and 0.2 mm in width, reddish brown in colour, long, with thick and smooth cuticle and bluntly conical posterior end, posterior end being truncated to form a prepuce-like fold. Head papilla present. Nerve ring 0.04 - 0.1 mm from anterior end. Oesophagus 2.3 - 2.6 mm in length.




Figure 1. Eustrongylides sp. (28X enlarged)

- a) Anterior part of the body
- (b) Posterior part of the body




Plate 1: Photomicrograph of *Eustrongylides* sp. (70X enlarged)

(a), (b) Anterior part of the body (c), (d) Posterior part of body.

Remarks: Due to the presence of the above characters the present nematode was considered as *Eustrongylides* sp. The present form was collected as larval form. Hence, it cannot identify upto species level.

Host: Channa punctata (New host record).

Habitat: Air bladder and intestine.

Locality: Karbhala wetland, Cachar District, Silchar, Assam (first geographical record)

Distribution of *Eustrongylides* in India: Bhopal; Maharashtra; Silchar, Assam.

Body of the present worm is long, reddish brown in colour, posterior end being truncated to form a prepuce-like fold. Head papilla presents in two circles. Oesophagus long. Due to the presence of the above characters the present nematode was considered as *Eustrongylides* sp.

CONCLUSION

The present worm is larval form of genus *Eustrongylides*, a nematode belonging to the family Dioctophymidae. During the present study, very less number of larval forms were collected from the fish host *Channa punctata* from Karbhala wetland, Assam. It is a new host record and new geographical record. Due to lack of mature structures, the present worm cannot identify upto species level.

ACKNOWLEDGEMENTS

The authors are thankful to the authorities of Assam Central University, Silchar and to the Principal and Head, Zoology Department, Thambal Marik College, Oinam, Manipur for giving the necessary laboratory facilities and to the UGC for providing a major research project to the second author and financial assistance to the first author. Thanks are due to Sri S. R. Dey Sarkar, retired Scientist, Zoological Survey of India, Kolkata; our teacher (Late) Prof. Umapati Sahay, retired Dean, HOD & Professor of Zoology, Ranchi University, Ranchi for helping in identification of specimens. Thanks are due to our teacher Prof. A. N. Jha, retired Professor, BRA Bihar University, Muzaffarpur for continuous advised and encouraging.

Note- This paper produced wherein facilities created under Biotech Hub, Department of Zoology, Thambal Marik College, Oinam sponsored by DBT, Ministry of Science and Technology, Government of India.

REFERENCES

- Jagerskiold, L. A. K. E. 1909. Zur Kenntnis der Nematoden-Gattungen Eustrongylides u. Hystrichis Soc. Acta. Ser. 4(2): 3-48.
- Canavan, W. P. N. 1929. Nematode parasites of vertebrates in the Philadelphia Zoological Garden and vicinity. *I. Parasit.* 21(1-2): 63-102.
- **3. Fernando C.H. and Furtado J. I. 1963.** A study of some helminth parasites of fresh water fishes in Ceylon. *Z.ParasitKde.*, **23:**141-163.
- **4. Kalyankar, S. D. 1974.** First report on *Eustrongylides* larva (Nematoda: Dioctophymidae) from fish in India. *Acta parasit. Pol.*, **22(30):** 331-333.
- 5. Naidu, T. S. V. 1979. A report on the larval forms of the nematodes belonging to *Gnathostoma* sp.; *Contracaecum* sp.; and *Eustrongylides* sp. from new piscian hosts, from Vidarbha region of Maharashtra state, India. *Current Science*. 48(5): 223-224.
- 6. Wang, W.; Li, L.; Yu, Y.; Feng, W.; Xiao, C.; Wang, G., Liu, J.; Yao, W.; Feng, S. 1997. Parasite fauna of fishes in Wuling Mountains area. In: Song D (ed) Invertebrates of Wuling mountains area. Science Press, Beijing, 73-146.
- Ibiwoye, T. I. I.; Balogun, A. M., Ogunsusi, R. A. and Agbontale, J. J. 2004. Determination of the infection densities of mudfish Eustrongylides in Clarias gariepinus and C. anguillaris from Bida floodplain of Nigeria. J Appl Sci Environ Manag, 8(2): 39-44.

- 8. Ibiwoye, T. I. I.; Okarme, A. N.; Ogunsusi, R. A. and Balogun, A. M. 2005. First report and record of nematode *Eustrongylides africanus* larvae in a vertebrate host mudfish *Clarias* species from Bida floodplain of Nigeria. *Afr. Sci.*, 6(2): 47-55.
- 9. Ibiwoye, T. I. I.; Ogunsusi, R. A.; Balogun, A. M. and Agbontale, J. J. 2008. The Contributions of environmental and haematological factors to the distributions and estimations of *Eustrongylides africanus* larvae densities in *Clarias gariepinus* and *Clarias anguillaris* from Bida floodplain of Nigeria. *J Fish Aquat Sci*, 3: 195-205.
- Bruce N. L. 1990. Hysterothylacium Ward and Magath, 1917, and Ichthyascaris Wu, 1949, ascaridoid nematodes from Australian demersal fishes. Memoirs of the Oueensland Museum 28: 389-426.
- Bruce N. L., Adlard R. D. and Cannon L. R. G. 1994.
 Synoptic checklist of ascaridoid parasites (Nematoda) from fish hosts. *Invertebrate Taxonomy*. 8: 583-674.
- **12.** Chabaud, A. G. 1965. Ordre des Ascaridida, *in* GRASSE P.-P. (ed.), *Traite de Zoologie*, tome IV, fascicule 3. Masson et Cie, Paris: 932-1025.
- **13.** Yamaguti, S. 1935. Studies on the helminth fauna of Japan. Pt. 9. Nematodes of fishes. *Jap. J. Zool.* **6(2)**: 337-386

Biospectra: Vol. 20(1), March, 2025

An International Biannual Refereed Journal of Life Sciences